


Microbial ecosystems of Antarctica

WARWICK F. VINCENT

*Taupo Research Laboratory,
Division of Marine and Freshwater Science,
Department of Scientific and Industrial Research,
New Zealand.*



CAMBRIDGE UNIVERSITY PRESS

Cambridge

New York New Rochelle

Melbourne Sydney

Published by the Press Syndicate of the University of Cambridge
The Pitt Building, Trumpington Street, Cambridge CB2 1RP
32 East 57th Street, New York NY 10022, USA
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© Cambridge University Press 1988

First published 1988

Printed in Great Britain at the University Press, Cambridge

British Library cataloguing in publication data

Vincent, Warwick F.

Microbial ecosystems of Antarctica.

1. Antarctica. Ecosystems

I. Title II. Series

574.5'0998'9

ISBN 0-521-32875-6

Library of Congress cataloguing in publication data

Vincent, Warwick F.

Microbial ecosystems of Antarctica.

(Studies in polar research)

Includes index.

1. Microbial ecology—Antarctic regions.

I. Title. II. Series.

QR100.V56 1988 576'.15'09989 88-7316

ISBN 0-521-32875-6

PN

Cover: Scanning electron micrograph of a cyanobacterial mat from Fryxell Stream, southern Victoria Land

Contents

Preface	xi
1 Introduction	1
2 Snow and ice ecosystems	6
2.1 Introduction	6
2.2 The environment	6
2.2.1 Snow environments	6
2.2.2 Glacier meltholes – the cryoconite environment	9
2.2.3 Ice-shelf systems	11
2.2.4 Ice-bubble habitats	17
2.3 Microbial community structure	18
2.3.1 Snow communities	18
2.3.2 Cryoconite communities	22
2.3.3 Ice-shelf communities	23
2.3.4 Ice-bubble communities	24
2.3.5 Continental ice-sheet flora	24
2.4 Microbial processes	25
2.5 Trophic structure	25
3 Sea-ice ecosystems	26
3.1 Introduction	26
3.2 The environment	27
3.2.1 Sea-ice cover	27
3.2.2 Ice formation and structure	29
3.2.3 Sea-ice chemistry	32
3.2.4 Light and temperature	33
3.3 Microbial community structure	34
3.3.1 Microalgal communities	35
3.3.2 Bacterial communities	44
3.3.3 Protozoan communities	46
3.3.4 Fungal communities	47

3.4	Microbial processes	47
3.4.1	Photosynthesis	47
3.4.2	Photoadaptation and nutrient deficiency	48
3.4.3	Heterotrophic production	49
3.4.4	Winter survival	51
3.5	Trophic structure	51
4	The marginal-ice zone	53
4.1	Introduction	53
4.2	The environment	54
4.2.1	Physical environment	56
4.2.2	Nutrients	57
4.3	Microbial community structure	59
4.3.1	Microalgal communities	59
4.3.2	Bacteria	63
4.3.3	Protozoa	63
4.4	Microbial processes	63
4.4.1	Ice-edge photosynthesis	63
4.4.2	Nitrate assimilation	66
4.4.3	Bacterial heterotrophy	67
4.4.4	Reproductive dynamics of protozoans	67
4.5	Trophic structure	67
5	The open ocean	70
5.1	Introduction	70
5.2	The environment	71
5.2.1	Bathymetry	71
5.2.2	Currents and mixing	71
5.2.3	Temperature	77
5.2.4	Light	77
5.2.5	Nutrients	78
5.3	Microbial communities	78
5.3.1	Microalgal communities	78
5.3.2	Bacterial communities	81
5.3.3	Protozoan communities	82
5.4	Microbial processes	83
5.4.1	Photosynthesis	83
5.4.2	Phototrophic production: the influence of light and temperature	85
5.4.3	Phototroph respiratory losses	87
5.4.4	Phototrophic nutrition	88
5.4.5	Bacterial heterotrophy	90
5.4.6	Protozoan grazing	93
5.5	Trophic structure	94

6 Benthic marine environments	97
6.1 Introduction	97
6.2 The environment	98
6.2.1 Environmental stability	98
6.2.2 Physical substrates	98
6.2.3 Sediment silica and organic carbon	100
6.2.4 Sediment nutrients	102
6.3 Microbial communities	102
6.3.1 Microalgal populations	102
6.3.2 Bacterial populations	103
6.3.3 Protozoan communities	104
6.4 Microbial processes	106
6.4.1 Photosynthesis	106
6.4.2 Bacterial heterotrophy	107
6.4.3 Protozoan physiology	108
6.5 Trophic structure	109
7 Lakes and streams	111
7.1 Introduction	111
7.2 The environment	111
7.2.1 Snow and ice cover	112
7.2.2 Salinity	119
7.2.3 Temperature	121
7.2.4 Stream discharge	122
7.2.5 Stream sediment load	124
7.3 Microbial communities	124
7.3.1 Phototrophs	125
7.3.2 Heterotrophic bacteria	131
7.3.3 Fungi	132
7.3.4 Protozoa	133
7.4 Microbial processes	135
7.4.1 Photosynthesis	135
7.4.2 Bacterial heterotrophy	139
7.4.3 Nutrient cycling	139
7.5 Trophic structure	148
8 Soil ecosystems	152
8.1 Introduction	152
8.2 The environment	153
8.2.1 Regional soil types	153
8.2.2 Moisture content	154
8.2.3 Temperature	156
8.2.4 Snow cover	157

8.2.5	Salinity	158
8.2.6	Soil pH	159
8.2.7	Organic content	160
8.2.8	Nitrogen content	161
8.3	Microbial community structure	162
8.3.1	Phototrophic communities	162
8.3.2	Bacterial heterotrophs	167
8.3.3	Fungal communities	169
8.3.4	Protozoan communities	171
8.4	Microbial processes	173
8.4.1	Photosynthetic production	173
8.4.2	Nitrogenase activity	173
8.4.3	Methanogenesis	174
8.4.4	Temperature–salinity adaptation	176
8.4.5	Microbial nitrogen transformations	176
8.4.6	Bacterial and fungal heterotrophy	177
8.5	Trophic structure	177
9	Lithic ecosystems: the rock environments	179
9.1	Introduction	179
9.2	The environment	179
9.2.1	Rock type	180
9.2.2	Moisture availability	182
9.2.3	Light availability (PAR)	182
9.2.4	Radiation and temperature	182
9.2.5	Nutrient supply	185
9.3	Microbial community structure	185
9.3.1	Phototrophic and fungal communities	185
9.3.2	Bacterial communities	191
9.4	Microbial processes	191
9.4.1	Phototrophic production	191
9.4.2	Biogenic exfoliation	193
9.5	Trophic structure	193
10	Microbial strategies in Antarctica	195
10.1	Introduction	195
10.2	Low temperatures	195
10.2.1	Cold avoidance	196
10.2.2	Cellular responses to chilling	196
10.2.3	Antarctic temperature response curves	201
10.2.4	Adaptation to chilling	207
10.2.5	Cellular freezing	208
10.2.6	Adaptation to freezing	211

10.3	High salinity	213
10.4	Dehydration	216
10.5	Low light	218
10.6	High radiation	220
10.7	Winter darkness	221
11	Microbes and humans in Antarctica	222
11.1	Introduction	222
11.2	Medical microbiology	224
11.3	Pollutants and microbial responses	226
11.4	Eutrophication in the Antarctic	228
11.5	Microbial dispersal	229
11.6	Endemism and displacement	231
	Glossary	233
	Appendix 1 Antarctic climates	238
1.1	Atmospheric pressure and average wind distribution	238
1.2	Precipitation and humidity	240
1.3	Radiation balance and air temperature	241
1.4	Climatic zones	246
1.5	Regional climate summaries	247
1.5.1	South Orkney Islands	247
1.5.2	Paradise Bay, Antarctic Peninsula	247
1.5.3	McMurdo Sound region	250
1.5.4	The 'Dry Valleys' region of southern Victoria Land	250
1.5.5	Vestfold Hills	252
1.5.6	Polar plateau	252
	Appendix 2 General environmental data	257
	References	265
	Index	291

Introduction

Extreme cold has shaped the antarctic environment and the microbial communities that live within it. Even throughout summer, air temperatures at the margins of the continent and in the maritime zone lie between -10 and $+5^{\circ}\text{C}$. The dramatic and discontinuous change in the physical properties of water over this range, in particular at its freezing point (Fig. 1.1), has a far-reaching impact on the chemical and physical characteristics of all potential habitats throughout the region. These environmental effects restrict the types of organisms that can be supported, and severely limit the timing and intensity of biological processes.

The freezing and melt cycle exerts the dominant influence on antarctic life forms, although not always directly. Persistent snow and ice are a feature of many antarctic environments (most obviously the continental ice sheet, and the vast expanse of sea ice that encircles the continent each year), even during the period of maximum biological activity. Freezing and melting dictate the areal extent of these environments that in turn modify other properties of the region – for example the distribution of water masses in the Southern Ocean (see Chapters 4, 5) and the regional variations in climate (Appendix 1). For the organisms which live within snow and glacier ice (see Chapter 2), the various forms of sea ice (see Chapter 3), or the permafrost-influenced soils (see Chapter 8), growth and reproduction are totally dependent upon the minute pockets of liquid water, and during freezing and melting these are subject to major fluctuations in volume, osmolarity, pH and temperature. Ice formation can physically disrupt environments (e.g. parts of the marine benthos subject to ice-scouring or anchor ice effects (see Chapter 6)) or cause large-scale changes in the salinity and gas content of the remaining water and the amount of light that penetrates down into it (e.g. ice-capped lakes and pools (see Chapter 7)). Rapid shifts backwards and forwards across the freezing point may effectively sterilise certain environments or force the




Fig. 1.1. Even in January, the warmest month for biological activity, air temperatures throughout much of Antarctica lie in the range -10 to $+5$ °C (the shaded bars represent the temperature range for each region, from mean daily minima to mean daily maxima; full data in Appendix 1). The abrupt shift in the properties of H_2O across this range, illustrated here by heat capacity (after Franks, 1985), has a major influence on the distribution and activity of the biota.

micro-organisms into thermally more stable parts of the habitat, e.g. abiotic rock faces and the well developed microbial communities at depth beneath the rock surface (see Chapter 9). Conversely, for certain habitats the formation of an ice cover provides insulating protection from the outside environment, and may allow liquid water to persist despite sub-zero air temperatures.

The subsequent melting process often occurs rapidly. This causes an abrupt improvement in growth conditions that may amplify the effects of the strongly seasonal light regime on primary production and subsequent food chain processes. Most of the Southern Ocean for example, contains sparse concentrations of microalgae (phytoplankton) that have low photosynthetic rates (see Chapter 5). When the sea ice breaks up each year, the

lower salinity meltwater restricts vertical mixing and retains the phytoplankton under improved light conditions for photosynthesis; this is believed to cause a band of intense primary and secondary production that sweeps across the ocean behind the retreating ice pack (see Chapter 4). Ice shelf (see Chapter 2) and meltwater stream (see Chapter 7) ecosystems in Antarctica are fundamentally linked to radiation and temperature, and small shifts in the energy balance at their ice faces may markedly alter the availability of water for aquatic life in these ecosystems. With summer air temperatures typically remaining close to 0 °C in many antarctic habitats, small changes in local climate can completely shift the balance between freezing and melting and thereby dramatically change the chemical and physical environment. As a result many antarctic ecosystems might be expected to show large interannual variations in the timing and duration of conditions favourable for biological processes.

Marked changes also take place in the immediate cellular environment during the freezing and melt cycle. Extracellular ice formation results in severe dehydration, often compounded by osmotic stress; intracellular ice formation generally causes the death of the cell. Antarctic microorganisms, like those in other freezing habitats have a range of adaptive strategies that allow them to avoid or at least minimise these destructive effects (see Chapter 10).

Cold temperatures above freezing may also exert a decisive influence on the activity and species composition of antarctic communities. At a cellular level these effects operate in part through the Arrhenius relationship between chemical reaction rate and temperature. However for many species in Antarctica and elsewhere there are major departures from this simple function, even within the ambient temperature range. Some of these responses to cold operate at the cellular macromolecule level. The extent of hydrogen-bonding of proteins with water molecules is a determinant of their catalytic activity and this surrounding water structure is sensitive to temperature. Similarly, temperature has a major influence on the fluidity of lipid membranes that in turn regulates many cellular functions (see Chapter 10). These biochemical and biophysical stresses may be exacerbated by further chemical interactions in habitats where freezing has generated high solute concentrations and unusual pH regimes.

Low temperatures may have a differential effect on select components of the community. In microbial consortia of phototrophs and heterotrophs, the markedly different responses to temperature by each functional component (e.g. fungal respiration versus net photosynthesis by the algae in the microscopic lichens that live within certain rocks (see Chapter 5)) can

further generate pronounced non-linearities in the temperature response curve. These effects of low temperature may strongly interact with other environmental variables including light intensity (e.g. net carbon fixation by the rock microbial communities), daylength (e.g. net photosynthesis in the Southern Ocean) and pH (e.g. methanogenesis in moss tundra soils). They may also influence the usefulness of traditional descriptors of microbial activity – for example chlorophyll *a* does not seem to decompose completely in various cold water environments and may be a misleading guide to phototrophic biomass in communities such as the marine benthos (see Chapter 6) and stream microbial mats (see Chapter 7).

Antarctic microbial communities are fiercely regulated by their extreme environment, but they may also have a reciprocal influence on the physical and chemical properties of their surroundings. For example, the presence of dense growths of diatoms in the sea ice substantially lessens its physical strength (Buinitsky, 1977) and may thereby hasten its decay. The communities within the rocks can accelerate the rate of erosion by exfoliation; these rock-dwelling microbes appear to live in a 'precarious equilibrium' between slow growth and destructive biotic and abiotic weathering (Friedmann & Weed, 1987). Similarly the growth of snow algae and ice mats can markedly alter the albedo of these environments and accelerate the rate of ablation. Dense populations of planktonic microalgae within ice-covered lakes can affect the distribution of heat and encourage relatively fast rates of mixing by penetrative convection (Matthews & Heaney, 1987).

Throughout Antarctica micro-organisms such as algae, bacteria, fungi and protozoa play a leading role in the biological transfer of materials and energy. In the extreme portions of this global region, particularly those experiencing near-continuous freezing or very rapid freeze–thaw cycles, higher plants and animals are completely absent and the food chains and biochemical nutrient cycles are entirely microbial. In many antarctic environments, however, human beings and the materials and microbes that we bring with us have generated new, additional tests of microbial resilience (see Chapter 11).

Perhaps the greatest challenge that microscopic communities in Antarctica may face will be the environmental shifts induced by perturbations of the Earth's atmosphere. Current climatic models that examine the effects of man-induced increases in atmospheric CO₂ and other gases predict a 0.5–2 °C warming of the temperate zone, and a more pronounced warming in the polar regions, by the middle of the next century (e.g. Schlesinger, 1986). An increase of the ground level temperatures in Antarctica by only a few degrees Celsius would alter the seasonal balance of freezing and

melting, with potentially greater ecological repercussions than elsewhere in the biosphere. Human influences on the ozone layer and the resultant increase in ultraviolet radiation particularly in the Antarctic (e.g. Farman, Gardiner & Shanklin, 1985) may also induce strong responses at the microbial level, and cause further shifts in genetic diversity.

True endemism appears to be relatively rare amongst the antarctic microbiota, but the environments of this region have selected for a wide range of robust microbial assemblages. The following chapters explore the structure of these assemblages, the chemical and physical properties of their surroundings and the interactions at both a population and cellular level between the microbiota and their antarctic habitats.